



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SOLUTION BY THE PROPOSER.

It appears that x, y, v, w , and a are to be rational numbers and $a \neq 0$.

If $v^2 = y^4$, $w^2 = x^4$, and a solution is given by $v = \pm y^2$ $w = \pm x^2$.

If $v^2 \neq y^4$ we may proceed as follows: From the given equation, we have

$$x^4 - w^2 = a(v^2 - y^4); \quad (2)$$

whence,

$$\frac{x^4 - w^2}{v^2 - y^4} = a = \frac{(x^4 - w^2)(v^2 - y^4)}{(v^2 - y^4)^2}, \quad (3)$$

or

$$\left(\frac{x^2 v \pm w y^2}{v^2 - y^4} \right)^2 - \left(\frac{x^2 y^2 \pm w v}{v^2 - y^4} \right)^2 = a. \quad (4)$$

Letting

$$\frac{x^2 v \pm w y^2}{v^2 - y^4} = b, \quad (5)$$

$$\frac{x^2 y^2 \pm w v}{v^2 - y^4} = c \quad (6)$$

and solving (5) and (6) for x^2 and w ,

$$x^2 = bv - cy^2 \quad (7)$$

and

$$w = cv - by^2. \quad (8)$$

From (7),

$$v = \frac{x^2 + cy^2}{b}, \quad (9)$$

Substituting the value of v from (9) in (8), we have

$$w = \frac{cx^2 + c^2 y^2 - b^2 y^2}{b}.$$

But from (4), (5), and (6), $b^2 - c^2 = a$. Hence if we put $b - c = m$, $b + c = a/m$; then

$$b = \frac{1}{2} \left(\frac{a}{m} + m \right), \quad c = \frac{1}{2} \left(\frac{a}{m} - m \right).$$

Hence

$$\begin{aligned} x &= r, & y &= s, \\ v &= \frac{2r^2 + \left(\frac{a}{m} - m \right) s^2}{\frac{a}{m} + m} = \frac{2mr^2 + (a - m^2)s^2}{a + m^2}, \\ w &= \frac{\left(\frac{a}{m} - m \right) r^2 - 2as^2}{\frac{a}{m} + m} = \frac{(a - m^2)r^2 - 2ams^2}{a + m^2}. \end{aligned}$$

2765 [1919, 171]. Proposed by A. M. HARDING, University of Arkansas.

ABC is an equilateral triangle. A point D is taken in BC such that BD is $\frac{1}{3}$ of BC and E is taken in CA such that CE is $\frac{1}{3}$ of CA . If the lines AD and BE intersect at O , show that OC is perpendicular to AD .

SOLUTION BY THE LATE L. G. WELD.

Since CD is twice CE and $\angle DCE = 60^\circ$ the auxiliary line DE is perpendicular to CE ; whence the point E is in the circumference of a circle described upon CD as a diameter. Since the triangles BCE and BOD are similar

$$BD \cdot BC = BO \cdot BE.$$

Hence, O , as well as E , lies in the circumference of the above circle and the angle COD , being inscribed in a semicircle, is a right angle.